

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
3 March 2005 (03.03.2005)

PCT

(10) International Publication Number
WO 2005/019470 A2

(51) International Patent Classification⁷: **C12Q**

(21) International Application Number:
PCT/US2004/025945

(22) International Filing Date: 10 August 2004 (10.08.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/494,482 11 August 2003 (11.08.2003) US
60/508,034 1 October 2003 (01.10.2003) US
60/512,941 20 October 2003 (20.10.2003) US

(71) Applicant (for all designated States except US): **ACLARA BIOSCIENCES, INC.** [US/US]; 1288 Pear Avenue, Mountain View, CA 94043 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHAN-HUI, Po-Ying** [CA/US]; 13056 Parkhurst Drive, Oakland, CA

94619 (US). **SINGH, Sharat** [US/US]; 27359 Julietta Lane, Los Altos, CA 94022 (US). **SALIMI-MOOSAVI, Hossein** [CA/US]; 1232 Vicente Drive, Sunnyvale, CA 94086 (US).


(74) Agent: **MACEVICZ, Stephen, C.**; Aclara Biosciences Inc., 1288 Pear Avenue, Mountain View, CA 94043 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: DETECTING AND PROFILING MOLECULAR COMPLEXES



(57) Abstract: Methods are provided for detecting the formation of complexes of molecules, especially proteins, in a sample, such as a cell or tissue lysate. In one aspect, a cleaving probe specific for a first protein in a complex and one or more binding compounds specific for one or more second proteins in a complex are provided. Upon binding, the cleaving probe is induced to generate an active species, such as singlet oxygen, that cleaves molecular tags attached to the binding compounds only in the local region of the cleaving probe. The released molecular tags are separated from the assay mixture and from one another to provide a readout that is related to the number and types of proteins present in the complex.



GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.